

<b>Disease</b>	Breast Cancer
<b>Regimen Name</b>	Dose-Dense AC (DOXOrubicin/Cyclophosphamide) followed by Paclitaxel
<b>Course</b>	Dose-Dense AC (DOXOrubicin/Cyclophosphamide) followed by Paclitaxel
<b>Cycle Length</b>	14-day cycle for 4 cycles
<b>Premedications</b>	The Emetic Risk: Day 1 High ( <b>start all before chemotherapy</b> )  <p style="text-align: center;"><b>Aprepitant</b> 125 mg PO once on Day 1</p> <p style="text-align: center;"><b>Palonosetron</b> 0.25 mg IV once on Day 1</p> <p style="text-align: center;"><b>Dexamethasone</b> 12 mg PO/IV once on Day 1</p>
<b>Regimen</b>	<b>DOXOrubicin</b> 60 mg/m <sup>2</sup> IV Push on Day 1 (Special instructions – not included at this time)  <b>Cyclophosphamide</b> 600 mg/m <sup>2</sup> IV over 30 minutes on Day 1 (special instructions-not included at this time)
<b>Fever Neutropenia</b>	High (Myeloid growth factors)  <p style="text-align: center;"><b>Filgrastim</b> 5 mcg/kg subcutaneously daily recommended to start 24 – 72 hours after completion of and to continue until post-nadir ANC recovery to normal or near normal levels by laboratory standards.</p>